

Volume 26 | Issue 2

Article 28

Rapid screening of toxic glycoalkaloids and micronutrients in edible nightshades (Solanum spp.)

Follow this and additional works at: https://www.jfda-online.com/journal

Part of the Food Science Commons, Medicinal Chemistry and Pharmaceutics Commons, Pharmacology Commons, and the Toxicology Commons



This work is licensed under a Creative Commons Attribution-Noncommercial-No Derivative Works 4.0 License.

Recommended Citation

Yuan, B.; Byrnes, D.; Giurleo, D.; Villani, T.; Simon, J.E.; and Wu, Q. (2018) "Rapid screening of toxic glycoalkaloids and micronutrients in edible nightshades (Solanum spp.)," *Journal of Food and Drug Analysis*: Vol. 26 : Iss. 2, Article 28. Available at: https://doi.org/10.1016/j.jfda.2017.10.005

This Original Article is brought to you for free and open access by Journal of Food and Drug Analysis. It has been accepted for inclusion in Journal of Food and Drug Analysis by an authorized editor of Journal of Food and Drug Analysis.

Available online at www.sciencedirect.com

ScienceDirect





Original Article

Rapid screening of toxic glycoalkaloids and micronutrients in edible nightshades (Solanum spp.)



Bo Yuan^a, David Byrnes^b, Daniel Giurleo^{b,c}, Thomas Villani^{b,c}, James E. Simon ^{b,c,*}, Qingli Wu ^{a,b,c,**}

^a Department of Food Science, 65 Dudley Road, New Brunswick, NJ 08901, USA

^b New Use Agriculture and Natural Plant Products Program, Department of Plant Biology, Rutgers University, 59 Dudley Road, New Brunswick, NJ 08901, USA

^c Department of Medicinal Chemistry, Ernest Mario School of Pharmacy, Rutgers University, 160 Frelinghuysen Road, Piscataway, NJ 08854, USA

ARTICLE INFO

Article history: Received 2 July 2017 Received in revised form 28 September 2017 Accepted 14 October 2017 Available online 11 November 2017

Keywords: Alkaloids Flavonoids Carotene Vitamin E Antioxidant activity Saponins Solanum

ABSTRACT

African indigenous vegetables (AIVs) because of their nutrient density have the unique potential to reduce micronutrient deficiencies in sub-Saharan Africa, yet some may also contain anti-nutritive compounds. Vegetable nightshades from Solanum americanum, Solanum nigrum, Solanum scabrum and Solanum villosum are among the major AIVs used as a leafy vegetables and consumed regularly in many countries in sub-Sahara Africa. These under-recognized food crops have not been subjected to extensive studies for their nutritional and antinutritive factors. In this study, 15 entries of the vegetable nightshades were field-grown and the leaves which are the consumed product of commerce chemically profiled by LC/ESI-MS. Twenty-three flavones, eight saponins, and two glycoalkaloids along with a phenolic acid of chlorogenic acid were identified by MS and UV data. Anti-nutrient glycoalkaloids were quantified as total aglycones after acidic hydrolysis using MS detection and found to be within safe-consumption thresholds by comparison with the glycoalkaloid level in the globally consumed Solanum member eggplants. Edible nightshades were also found to be sources of β -carotene, vitamin E and total polyphenols and exhibited high antioxidant activity. Results of this study support that consumption of vegetable nightshades are safe from the presence of glycoalkaloids and thus, can contribute to the reduction of micronutrient deficiency in sub-Sahara Africa.

Copyright © 2017, Food and Drug Administration, Taiwan. Published by Elsevier Taiwan LLC. This is an open access article under the CC BY-NC-ND license (http:// creativecommons.org/licenses/by-nc-nd/4.0/).

E-mail addresses: jimsimon@rutgers.edu (J.E. Simon), qlwu@sebs.rutgers.edu (Q. Wu). https://doi.org/10.1016/j.jfda.2017.10.005

^{*} Corresponding author. Department of Plant Biology, Rutgers University, 59 Dudley Road, New Brunswick, NJ 08901, USA.

^{**} Corresponding author. Department of Plant Biology, Rutgers University, 59 Dudley Road, New Brunswick, NJ 08901, USA.

^{1021-9498/}Copyright © 2017, Food and Drug Administration, Taiwan. Published by Elsevier Taiwan LLC. This is an open access article under the CC BY-NC-ND license (http://creativecommons.org/licenses/by-nc-nd/4.0/).

1. Introduction

African indigenous vegetables (AIVs) are consumed as important nutrient-rich foods locally and regionally in sub-Saharan Africa, with many also utilized for their medicinal properties [1]. Such AIVs, also called traditional African vegetables, are collected from the wild or cultivated to a limited extent and consumed or marketed, serving as an important income generating opportunity for the typical small-scale farmer, especially in such economically limited regions [2]. Adapted to the local environment, AIVs often provide more sustainable production than exotic or introduced crops such as European vegetables [3]. Efforts are being made to increase the farming and marketing of AIVs in an attempt to alleviate hunger and improve nutrition, and to increase farmers income, improving the local and regional economy [3].

African nightshades are among the most popular and as such high priority leafy AIVs, and their leaves as well as tender stems can be steamed and eaten like spinach and amaranth. The edible nightshades represent a wide group of botanically and genetically related plants belonging to approximately 30 species in the Solanum genus of the Solanaceae family, and are diversely referred to as vegetable nightshades, edible nightshades, garden nightshades, common nightshades, 'S. *nigrum* complex', or 'S. *nigrum*' and related species [4]. Despite their frequently reported nutritional attributes, Solanum species are also well known to contain toxic alkaloids, such as glycosides of solasodine and solanidine [5]. This safety concern is associated with the edible African Solanum nightshade species, as these compounds are known to be present in the fruits [6] and have limited the promotion of their cultivation and marketing.

As edible nightshades are consumed in sub-Sahara Africa for their leaves, not the fruits, the presence of glycoalkaloids in leaves could present health concerns. Therefore, the purpose of this research was to determine the nutritional content of edible African nightshades and to also examine whether glycoalkaloids are present in the leaves, and if so to identify and quantitate such compounds. The results would therefore clarify consumption safety concern and promote the cultivation and marketing of leafy AIVs, or identify if any sources are free and devoid of such compounds for breeding and crop improvement programs. In this investigation, 15 African edible nightshade entries were chemically profiled by LC/UV/ MS to identify major bioactive compounds as well as toxic glycoalkaloids. Additionally, the other nutritive values were measured including β -carotene and vitamin E level, total polyphenol content (TPP) and total antioxidant activity.

2. Materials and methods

2.1. Chemical reagents

Standard compounds solasodine was purchased from MP Biomedicals (Santa Ana, CA, USA) and solamargine from MedChem Express (Monmouth Junction, NJ, USA). Standards β -carotene and vitamin E (α -tocopherol), Folin Ciocalteu's phenol reagent, 6-hydroxy-2,5,7,8-tetramethylchromane-2-

carboxylic acid (Trolox) and 2,2'-azino-bis (3-ethylbenzothiazoline-6-sulphonic acid) (ABTS) from Sigma-–Aldrich (St. Louis, MO, USA). Gallic acid was purchased from Acros Organics (Belgium, WI, USA) and acetone from BDH Chemicals (Radnor, PA, USA). Methanol, ethyl acetate, *tert*butyl methyl ether, concentrated hydrochloric acid, and HPLC grade water and acetonitrile modified with 0.1% formic acid were purchased from Fisher Scientific (Fair Lawn, NJ, USA).

2.2. Equipment

A propane-heated walk-in Powell Maxi Miser tobacco dryer (Bennettsville, SC) was used for sample drying. Agilent 1100 series LC/MSD instrument (Waldbronn, Germany) was used for phytochemical profiling. The HPLC was equipped with an auto-degasser, quaternary pump, thermostatted column compartment and a diode-array detector (DAD). Column Agilent Polaris 3 Amide C18, 250×4.6 mm (Santa Clara, CA, USA) was used for phytochemical profiling, and column Phenomenex Prodigy ODS-3150 \times 4.6 mm, 5 μm (Torrance, CA, USA) was used for quantification of total glycoalkaloids. The HPLC-MS interface used an electrospray ionization source (ESI) and the MS featured an ion trap analyzer. The software used was HP ChemStation, Bruker Daltonics 4.1 and DataAnalysis 4.1. Waters 2695 HPLC (Milford, MA, USA) was used for β-carotene and vitamin E measurement, which was equipped with a quaternary pump and a DAD. The separation was achieved by YMC-C30 carotenoid column, 5 μ m, 250 \times 4.6 mm (YMC Co., Ltd). The software was Millennium 4.00. Bio-Tek Synergy HT Multi-Mode Microplate reader (Winooski, VT, USA) was used for spectrophotometric measurement for total polyphenol assay and antioxidant assay. The software used was Bio-Tek KC4 Version 3.4.

2.3. Plant samples

Seeds of 15 entries of S. nigrum, Solanum scabrum, Solanum americanum and Solanum villosum (Table 3) were sown under greenhouse conditions at the Rutgers Research Greenhouses in New Brunswick, NJ. After four weeks of growth, the seedlings were transplanted during the first week of June in 2015 into a cultivated field at the Clifford E. & Melda C. Snyder Research and Extension Farm, New Jersey Agricultural Experiment Station of Rutgers University in Pittstown, New Jersey (40.6°N, 75.0°W, 116 m elevation). The leaves of the nightshades were manually harvested with the first harvest occurring 21–28 days post-field transplanting. The aerial parts were cut ~15 cm above the soil line to allow the plants to regrow for multiple harvesting.

2.4. Sample preparation

The collected aerial parts were dried at 40 C° for two weeks and then ground into powder. The samples were stored in shaded zip-lock bags under room temperature. For phytochemical profiling by LC/UV/MS, total polyphenol (TPP) assay and antioxidant assay, around 200 mg of the sample was accurately weighed and extracted with 25 mL 70% methanol with 0.1% formic acid. Each extract was fully vortexed, sonicated in a water bath for 5 min, and then let stand still

Table 1 – Peak assig	nments used :	for the analysis of 15 edible nightshade	e (Solanum spp.) accessi	ons.		
Peak/Compound code	RT(min)	Compound ID	[M+H] ⁺ /[M+Na] ⁺ (m/z)	Fragment ion (m/z)		
1	13.6	Chlorogenic acid	377, 355 ^ª	_		
2	14.0	Ouercetin-G-Rha-G	795 ^a	611, 465, 303		
3	16.2	Quercetin-G-Xyl-Xyl	751 ^a	597, 465, 303		
4	16.8	Quercetin-G-Rha-Rha	779 ^a	611, 465, 303		
5	17.8	Quercetin-G-G	649 ^a	465, 303		
6	18.1	Quercetin-G-Rha-Xyl	765 ^a	465, 303		
7	18.3	Solasodine-G-G-Rha	884	576, 414		
8	18.7	Quercetin-G-Xyl	619 ^a	465, 303		
9	18.9	Quercetin-G-Xyl	619 ^a	465, 303		
10	19.1	Kaempferol-G-G	633 ^a	449, 287		
11	20.0	Solasodine-G-Rha-Rha	868	722, 576, 414		
12	20.2	Kaempferol-G-Xyl	603 ^a	449, 287		
13	20.6	Kaempferol-G-Xyl	603 ^a	287		
14	20.7	Kaempferol-G-G	633 ^a	449, 287		
15	21.4	Rhamnetin-G-Xyl-Xyl	765 ^a	611, 479, 317		
16	21.9	Quercetin-G-Rha	633ª	465, 303		
17	24.0	Quercetin-G	487 ^a	303		
18	24.0	Rhamnetin-G-G	663 ^a	479, 317		
19	25.3	Rhamnetin-G-Xyl	633ª	479, 317		
20	25.5	Kaempferol-G-Xyl	617 ^a	449, 287		
21	25.6	Rhamnetin-G-Xyl	633ª	479, 317		
22	26.9	Rhamnetin-G-Xyl	633ª	479, 317		
23	27.4	Kaempferide-G-Xyl	617 ^a	301		
24	27.8	Kaempferide-G-Xyl	617 ^a	463, 301		
25	28.6	Rhamnetin-G-Xyl	633 ^a	479, 317		
26	29.5	Rhamnetin-G-Rha	647 ^a	479, 317		
27	33.3	Tigogenin-G-G-G-G	1249 ^a	1087, 903, 741, 579, 417		
28	34.4	Tigogenin-G-G-G-Xyl-G	1197	1057, 903, 741, 579, 417 ^a		
29	34.8	Dehydrodiosgenin-G-G-Rha-Rha	1029	883, 737, 575, 413		
30	35.7	Diosgenin-G-G-Rha-Rha	1031	885, 739, 577, 415		
31	41.5	Tigogenin-G-G-Rha-Xyl-Xyl	1173 ^a	887, 741, 579, 417		
32	42.6	Tigogenin-G-G-Rha-Xyl-Xyl	1173 ^a	887, 741, 579, 417		
33	43.1	Dihydromethyldiosgein-G-Rha-Rha-G	1069 ^a	885, 739, 593, 431		
34	45.6	Tigogenin-G-G	903	741, 579, 417		
^a Sodium adducted ion	s: RT. retention t	ime: G glucosyl galactosyl mannosyl or othe	r hexosyl: Rha_rhamnosyl:	Xvl. Xvlosvl.		

overnight under room temperature [7–9]. The extract was centrifuged at 3000 rpm/min for 10 min. For TPP assay, the supernatant was directly ready for analysis. For antioxidant activity assessment, the supernatant was diluted twice by 70% methanol with 0.1% formic acid before the assay. For phytochemical screening by LC/ESI-MS, the supernatant was filtered by 0.45 µm filter prior to HPLC injection.

For quantification of total alkaloid aglycone, the sample was prepared based on protocols by P.G. Crabbe [10] with modification. Briefly, around 400 mg of sample was accurately weighed and hydrolyzed by 20 mL 0.5 M anhydrous methanolic hydrochloric acid in an eight dram vial with a screw cap, and incubated in a 70 °C water bath for 120 min. After incubation, the hydrolyte was cooled down and basified with 3 mL saturated sodium hydroxide methanolic solution to terminate the hydrolysis reaction, and was then brought to 25 mL by methanol. The final hydrolyte was filtered by 0.45 µm filter before LC/MS injection. For construction of the calibration curve and determination of limit of quantification (LOQ) and limit of detection (LOD), 4.7 mg solasodine standard was dissolved in 10 mL methanol as stock solution, and diluted 100 times as the first work solution, followed by successive two-fold dilution down to sub-nanogram/mL. LOQ was defined as the signal to noise ratio (S/N) at 10:1, and LOD as S/N at 3:1. The recovery rate was evaluated by spiking known amount of authentic standard solamargine to approximately 200%, 100% and 50% of the expected values in the sample in terms of total aglycone and then hydrolyzing following the same procedure.

The aforementioned hydrolysis method was also applied to release aglycones of flavonoids and saponins for further confirmation of their structures.

For the analysis of β -carotene and vitamin E, about 500 mg sample was weighted accurately and extracted by 5 mL acetone in a two dram vial, fully vortexed and sonicated for 30 s. The extract was then centrifuged at 3000 rpm for 5 min with the supernatant decanted to an 8 dram amber vial. The sample was subsequently extracted in like manner by another 5 mL acetone and 2 mL tert-butyl methyl ether, respectively, with the supernatants combined in the 8 dram vial. The supernatant was then rotovapped to yield the dry extract. The dried residue was then reconstituted into 2 mL 1:1 ethyl acetate: methanol, followed by centrifugation at 3000 rpm for 10 min, and the supernatant was ready for HPLC analysis. The extract for each entry was prepared in triplicate.

Table 2 – Phytochemical profile analyses of 15 edible nightshades (Solanum spp.) accessions.															
Peak no.	S.n 1	S.n 2	S.n 3	S.n 4	S.n 5	S.s 1	S.s 2	S.s 3	S.s 4	S.s 5	S.s 6	S.s 7	S.s 8	S.a 1	S.v 1
1	_	Т	+	Т	T	+	_	+	+	+	+	Т	Т	+	Т
2	_	_	_	_	+	+	_	+	+	Т		_	_	++	++
3	-	-	+	-	_	-	-	-	_	-	-	-	-	_	-
4	-	-	-	+	Т	$^{++}$	+	$^{++}$	+	+	++	+	+		-
5	$^{++}$	++	+	Т	++	+	Т	+	+	+	Т	Т	Т	++	++
6	-	-	-	-	_	-	Т	+	+	+	Т	-	-	_	-
7	-	-	-	-	-	-	-	-	_	-	-	-	-	-	Т
8	-	-	+	-	_	-	-	-	—	_	-	-	-	_	-
9	-	-	++	-	_	-	-	-	—	_	-	-	-	_	-
10	+	+	-	-	+	-	-	-	-	-	-	-	-	Т	-
11	-	-	-	-	Т	-	-	-	_	-	-	-	-	-	Т
12	-	-	Т	-	-	-	-	-	-	-	-	-	-	-	-
13	-	-	Т	-	-	-	-	-	-	-	-	-	-	Т	-
14	-	-	-	-	-	-	-	Т	-	Т	-	-	-	-	-
15	-	-	++	-	-	-	-	-	-	-	-	-	-	-	-
16	Т	-	+	+	Т	+	+	++	+	+	++	+	+	-	Т
17	+	Т	+	-	+	-	-	-	-	-	-	-	-	Т	
18	-	-	+	-	Т	-	-	-	-	-	-	-	-	++	+
19	-	-	++	-		-	-	-	-	-	-	-	-	-	-
20	-	-	-	Т		Т	-	Т	Т	Т	Т				-
21	-	-	++	-	-	-	-	-	-	-	-	-	-	-	-
22	-	-	Т	-	_	-	-	-	—	-	-	-	-	-	-
23	-	-	Т	-	_	-	-	-	—	-	-	-	-	-	-
24	-	-	Т	-	-	-	-	-	-	-	-	-	-	-	-
25	-	-	Т	-	—	-	-	-	—	-	-	-	-	-	-
26	-	-	+	-	—	-	-	-	—	-	-	-	-	-	-
27	+	+	-	+	+	+	+	+	+	+	+	+	+	+	+
28	++	++	-	++	++	++	++	++	++	++	++	++	++	++	++
29	-	-	++	-	-	-	-	-	-	-		-	-	-	-
30	-	-	++	-	-	-	-	-	-	-		-	-	-	-
31	+	++	-	+	++	++	+	++	++	++	+	+	+	++	+
32	+	+	-	+	++	++	+	++	++	++	+	+	+	+	+
33	-	-	++	-	-	-	-	-	-	-	-	-	-	-	-
34	+	++	-	-	-	-	-	-	-	-	-	-	-	-	+

Compound codes refer to Table 1 and the plant sample code are the same as in Table 3. "++", strong peaks defined as intensity of UV–vis over 10 mAU or of extracted ion chromatograph (EIC) by extraction of molecular ions and corresponding fragment ions over 5×10^5 ; "T", trace level, defined as intensity of UV–vis lower than 1 mAU or EIC intensity lower than 5×10^4 ; "+", peak intensity between "++" and "T"; "-", peaks not detected.

2.5. Phytochemical profiling

The aerial part extract was chemically profiled by HPLC/UV/ ESI-MS. HPLC grade water with 0.1% formic acid was used for mobile phase A and acetonitrile with 0.1% formic acid for mobile phase B. The gradient started from 10% to 20% B in 0-10 min, 20% to 28% in 10-30 min, 28%-30% in 30-40 min, and 30% to 50% in 40-60 min, followed by 5 min column flushing with 80% B and another 5 min column equilibration with the starting mobile phase before the next injection. The flow rate was 1 mL/min and the injection volume was 10 μ l. The column was kept at 25 °C. The DAD detector was set at 210 nm, 254 nm, 280 nm and 370 nm for signals, and scanning range 200-550 nm with 2.0 nm step for spectrum. About a third of the HPLC eluent was split into the MS detector. In the ESI, the nebulizer needle voltage was set at 3500 V of positive polarity. High purity nitrogen was used as both nebulizing gas at 40 psi and drying gas at 350 °C with a flow rate of 10 L/min. High purity helium was used as the collision gas and the collision energy was set at arbitrary compound stability 80%. The scanning mode was set at positive and the range at 200-1300 m/z.

2.6. Quantification of total glycoalkaloids in hydrolyzed nightshade samples using LC/MS

For MS detection, selected ion monitoring (SIM) mode was used for the selected target of solasodine at 414.7 m/z with a band width of 1.0 m/z, and the HPLC gradient started from 20% to 40% in 10 min, followed by 5 min column flushing with 80% B. The column was equilibrated by the starting mobile phase for 5 min before each injection. The other HPLC/MS parameters remained the same as phytochemical profiling. The aglycone content was reported as $\mu g/g$ dry weight (DW).

2.7. β -carotene and vitamin E analysis

The β -carotene and vitamin E content was analyzed using Waters HPLC. Mobile phase A was micron-filtered 98:2 methanol:1 M ammonium acetate, and mobile phase B was HPLC grade ethyl acetate, and both were manually degassed by sonication under vacuum. The gradient started from 0% to 35% B in 0–8 min, 35% to 40% in 8–28 min, 40% to 50% in 28–32 min, and 50%–60% in 32–36 min and held until 40 min. The injection volume was 10 µl. UV chromatograms for β -

Table 3 – Identification, alkaloid aglycone and nutrients level from 15 accessions of edible nightshades (Solanum spp.).										
Sample code ^a	Source ^b	Source ID	Glycoalkaloid aglycone (μg/g DW)	Vitamin E (μg/g DW)	β-carotene (µg/g DW)	ABTS (TE mg/g DW)	TPP (GAE mg/g DW)			
S.n 1	Simlaw Seeds (Kenya)	-	ND	106.9 ± 2.0^{def}	78.4 ± 4.5^{bcd}	16.80 ± 0.64^{gh}	8.54 ± 0.33^{gef}			
S.n 2	USDA	PI 306400	ND	92.0 ± 2.6^{gf}	28.1 ± 1.1^{f}	$15.49 \pm 0.69^{\rm h}$	$7.25 \pm 0.36^{\rm h}$			
S.n 3	USDA	PI 312110	ND	229.7 ± 19.9^{a}	141.7 ± 11.7^{a}	23.45 ± 0.68^{abc}	11.92 ± 0.49^{ab}			
S.n 4	USDA	PI 381289	ND	125.9 ± 7.6 ^{def}	70.6 ± 2.5^{ced}	18.34 ± 0.77^{gf}	9.58 ± 0.33 ^{de}			
S.n 5	USDA	PI 381290	7.1 ± 2.3^{a}	197.1 ± 17.9^{ab}	86.7 ± 8.9^{bc}	23.93 ± 1.12^{ab}	13.07 ± 0.42^{a}			
S.s 1	WorldVeg	SS 52	ND	114.9 ± 8.6 ^{def}	87.4 ± 3.7 ^{bc}	21.36 ± 0.50^{dc}	10.8 ± 0.35^{bc}			
S.s 2	WorldVeg	Ex Hai	ND	90.4 ± 3.7 ^{gf}	64.8 ± 0.3^{ed}	17.92 ± 0.45 ^{gf}	7.98 ± 0.26^{ghf}			
S.s 3	WorldVeg	SS 49; Olevolosi	ND	141.3 ± 1.9 ^{de}	96.0 ± 2.8^{b}	22.46 ± 1.11^{bdc}	12.21 ± 0.53^{a}			
S.s 4	WorldVeg	SS 04.2	ND	64.1 ± 2.8^{g}	65.2 ± 2.0^{ed}	$18.11 \pm 0.62^{ m gf}$	9.39 ± 0.28^{de}			
S.s 5	WorldVeg	BG 16; Nduruma	ND	192.5 ± 28.5^{ab}	91.5 ± 13.0^{bc}	25.00 ± 0.70^{a}	12.47 ± 0.42^{a}			
S.s 6	WorldVeg	BG-29	ND	102.1 ± 11.7 ^{gef}	55.1 ± 7.3 ^e	19.14 ± 0.33^{ef}	9.13 ± 0.24^{def}			
S.s 7	USDA	Grif 14198	ND	121.1 ± 8.6 ^{def}	87.7 ± 4.7^{bc}	21.26 ± 0.59^{d}	$10.98 \pm 0.40^{\rm bc}$			
S.s 8	USDA	PI 643126	ND	183.4 ± 30.1 ^{bc}	87.2 ± 14.0^{bc}	16.22 ± 0.33^{gh}	7.75 ± 0.52^{gh}			
S.a 1	USDA	PI 268152	ND	145.5 ± 5.2 ^{dc}	95.8 ± 3.8^{b}	24.81 ± 0.09^{a}	12.44 ± 0.43^{a}			
S.v 1	USDA	Grif 16939	$38.0 \pm 11.4^{\rm b}$	$114.3 \pm 5.0^{\text{def}}$	$138.1\pm4.0^{\rm a}$	20.99 ± 0.99^{ed}	10.26 ± 0.40^{dc}			

The results were reported as mean \pm STD (n = 3). ^{a-h}, values with same letters in the same column are not significantly different (ANOVA with Tukey's HSD test, p < 0.05).

^a S.n: Solanum nigrum; S.s: S. scabrum; S.a: S. americanum; S.v: S. villosum. ND, not detected.

^b Seeds from USDA were maintained by Plant Genetic Resources Conservation Unit, Griffin, GA. Seeds from World Vegetable Center (WorldVeg) were maintained by the regional center of east and southern Africa, Arusha, Tanzania. Italicized values of β-carotene contents were identified to be a source of vitamin A based on retinol equivalent (RE), assuming 90% moisture content in fresh vegetables [24,25].

carotene and vitamin E were generated at their maximum peak absorption wavelength of 452 nm and 295 nm, respectively, and both compounds were identified in comparison with their retention time and UV–Vis spectrum of the authenticated standards. For the calibration curve, around 20 mg vitamin E and 10 mg β -carotene standards were accurately weighed and dissolved in 25 mL ethyl acetate as stock solutions, respectively [11]. The stock solutions were then diluted by ethyl acetate in series to generate work solutions of ~4–~150 µg/mL for both compounds. The β -carotene and vitamin E content in samples was reported as µg/g DW.

2.8. Total polyphenol (TPP)

The method of TPP assay was based on that proposed by Singleton [12] and Kao et al. [13] with modification. First, 50 mL Folin Ciocalteu's reagent was diluted by distilled water to 500 mL, then 900 μ L Folin reagent was mixed with 80 μ L sample extract followed by the addition of 400 μ L saturated sodium carbonate solution. The reaction system was fully vortexed and let stand still for 1 h. Then 200 µL supernatant was transferred to a 96 multiple well and subject to absorption measurement under 765 nm. As to the calibration curve, 20.0 mg gallic acid was dissolved in 5 mL 70% methanol as the stock solution, and diluted to a series of work solutions with concentrations ranging from 15.6 µg/mL to 250 µg/mL 200 µL 70% methanol was used as negative control. The assay for each calibrator and sample was triplicated. Total polyphenol content in samples was expressed as the amount of gallic acid equivalent (GAE)/g DW.

2.9. Antioxidant activity

The antioxidant activity assay was based on the method proposed by Re et al. [14] and Nagulsamy [15] with

modification. First, 31.7 mg ABTS and 8.6 mg potassium persulfate were dissolved in 10 mL water and let stand still in darkness under room temperature for 12–16 h to form stable radical, and diluted to an absorption of ~1.3 at 734 nm. Next, 200 μ l ABTS was mixed with 20 μ l sample extract and let stand still for 15 min under room temperature, and the absorption was measured at 734 nm. For the calibration curve, 12.5 mg standard Trolox was dissolved in 5 mL pure ethanol as the stock solution, and diluted to series of work solutions with concentrations ranging from 19.5 μ g/mL to 195.3 μ g/mL. The assay for each calibrator and sample was conducted in triplicate. Antioxidant activity was expressed as the amount of Trolox equivalent (TE)/g DW.

2.10. Statistics

Data were represented in the form of mean \pm the standard deviation (STD) of three replicates of each sample. Data was analyzed using student's t-test and one-way analysis of variance (ANOVA) followed by Tukey's Honest Significant Difference (HSD) test (p < 0.05). The relationship between TPP content and antioxidant activity was evaluated by Pearson's correlation. All statistics was performed using SAS University Edition (SAS Institute Inc., Cary, NC, USA).

3. Results and discussion

3.1. Phytochemical profiling

Major peaks were identified based on the retention time, UV–Vis spectrum, and MS spectral data, as well as by comparison with authentic standards. The identities, retention time and MS data of individual compounds were summarized in Table 1. In the four *Solanum* species, flavonoids and saponins dominate the secondary metabolites portfolio along with phenolic acid chlorogenic acid, occasionally with trace level of glycoalkaloids. Apparent variance in chemical profile was observed within species, as well as striking similarities between them, as commonly observed in the *Solanum* complex [16]. The chemical profile of 15 different accessions was summarized in Table 2.

3.1.1. Flavonoids and phenolic acid

Flavonoids were previously reported as an important class of bioactive compounds in nightshades [16]. In this work, a major phenolic acid chlorogenic acid along with an abundance of flavonols were detected and identified in the four AIVs, including glycosylated derivatives of quercetin, kaempferol, rhamnetin and kaempferide. Simultaneous UV and MS chromatograms of a representative nightshade extract were illustrated in Fig. 1. The individual peaks were identified based on UV-Vis spectrum and MS data. For the major phenolic acid, compound 1 (RT 13.6 min) exhibited adduct molecular ions at 377 ($[M+Na]^+$) and 355 m/z ($[M+H]^+$) and maximum UV peaks at 208 nm, 245 nm and 328 nm, suggesting a chlorogenic acid, neochlorogenic acid or cryptochlorogenic acid which differ at the esterification site of the quinoyl [17]. This compound was further confirmed to be chlorogenic acid by comparison with authentic standards. For identification of flavonols, for example, compound 8 (RT 18.7 min) featured a sodium adduct molecular ion at 619 m/z $([M+Na]^+)$, which was fragmented into 465 m/z ([M $xylosyl + H^+$, followed by aglycone ion at 303 m/z ([M-xylosylhexosyl + H]⁺) which corresponded to the molecular mass of protonated quercetin. The maximal UV—vis absorption was around 360 nm, in agreement with that of quercetin glycosides in literature [17]. Thus, compound **8** was identified as quercetin conjugated with hexosyl and xylosyl. The above identification principle also applied to the other flavonol glycosides. In addition, the identification of aglycones of quercetin, kaempferol and rhamnetin were further confirmed by comparison with retention time of authentic standards after acid-assisted hydrolysis.

3.1.2. Glycoalkaloids

Glycoalkaloids were previously reported in leaves of all four Solanum species, though sometimes in disagreement. S. nigrum was extensively reported to contain solasodine glycosides including solamargine and solasonine and those with varied oligosaccharide chain [18,19]. Solanine, a glycoside of solanidine, was only occasionally reported at a content of 0.043 mg/g [20]. In S. scabrum, solamargine and solasonine was reported to be present [21], while the existence of alkaloids was decided to be "doubtful" in another study [22]. In a recent research, a screening by LC-QToF-MS revealed a lack of glycoalkaloids in the methanol extracts of both S. scabrum and S. villosum [23]. S. americanum was reported to contain glycosides of solasodine [24].

In this study, no alkaloids were detected in the leaves of the plants except in two accessions, S. villosum Grif 16939 and S. nigrum PI 381290, where two glycosides of solasodine was found (Fig. 2A). Compound **11** (RT 20.0 min), for example, has a molecular ion peak at 868 m/z [M+H]⁺, followed by fragment ion 722 m/z [M-rhamnosyl + H]⁺ and 576 m/z [M-rhamnosyl



Fig. 1 – Representative HPLC/UV/MS chromatograms of Solanum nigrum PI 312110. (A) UV–vis chromatogram at 370 nm. (B) Processed MS chromatogram extracting molecular ions and characteristic fragment ions. The identities, retention time and MS of each peak are listed in Table 1.



Fig. 2 – Processed MS chromatograms showing glycoalkaloids in Solanum villosum Grif 16939 (A) and saponins in Solanum villosum Grif 16939 (B) and Solanum nigrum PI 312110 (C).

rhamnosyl + H]⁺, and finally 414 m/z [M-rhamnosyl-rhamnosyl-hexosyl + H]⁺ corresponding to the mass of aglycone solasodine (Fig. 3). Thus, compound **11** was identified as solasodine-hexosyl-rhamnosyl-rhamnosyl (solamargine), and the structure was further confirmed as solamargine by comparison with the retention time and mass spectrum of authentic standard. Compound **7** (RT 18.3 min) was identified in like manner and speculated to be solasonine (solasodine-

hexosyl- hexosyl-rhamnosyl) as among the most reported solasodine glycosides in the S. *nigrum* complex. The elution pattern and mass spectra of these two compounds are also in agreement with the literature [23].

3.1.3. Saponins

Saponin identification was primarily based on mass spectrum interpretation and literature review. Compound 28 (RT 34.4 min), for example, had a molecular ion peak at 1197 m/z[M+H]⁺, which occasionally featured adducted ions at 1219 m/z $[M+Na]^+$ and 1237 m/z $[M+Na+H_2O]^+$. The parent ion underwent a loss of hexosyl moiety to generate sodium adducted fragment ion 1057 m/z [M-hexosyl + Na]⁺, whose remaining saccharide unites were then successively cleaved off to generate the fragment ions at 903 m/z [M-hexosylxylosyl]⁺, 741 m/z [M-hexosyl-xylosyl -hexosyl + H]⁺, 579 m/z $[M-hexosyl-xylosyl-hexosyl-hexosyl + H]^+$, and finally the aglycone ion at 417 m/z [M-hexosyl-xylosyl-hexosyl-hexosylhexosyl + H]⁺, which corresponded to the mass of tigogenin as extensively found in Solanum complex [6,8,24]. Thus, compound 28 was identified as tigogenin conjugate with four hexosyls and one xylosyl. The other saponins were identified in similar manner. In addition, the structures of aglycones of diosgenin and tigogenin were further confirmed by comparison with authentic standards after acid-assisted hydrolysis. The representative chromatograms are shown in (Fig. 2B and C).

3.2. Quantification of total glycoalkaloid aglycone

As the toxicity of glycoalkaloids has long been the major concern in the consumption and promotion of *S. nigrum* complex, the glycoalkaloids were quantified as the top priority to shed light on the potential consumption risk. Given the potential existence of occasionally occurring solasodine



Fig. 3 – Mass spectrum of solamargine (compound 11 in Fig. 2A) with structure and MS fragment pathway.

glycosides with varied oligo/monosaccharide moieties below the detection limit, and that the aglycone part was the major source of glycoside bioactivity, in this work, the plant samples were hydrolyzed during extraction to yield the uniform aglycone solasodine as a reflection of glycoalkaloids content. Using the method developed here, the glycoalkaloid positive samples indicated in the phytochemical profiling were hydrolyzed and analyzed using MS detection under SIM mode. The calibration curve using solasodine standard was Y = 16,578X+2,96,833, $R^2 = 0.9943$, with a linearity range of 4.7 ng/mL ~47 µg/mL. The LOQ was 4.7 ng/mL, and LOD was 2.3 ng/mL. The recovery rate at ~50%, ~100% and ~200% level was 102.9 ± 25.8%, 81.5 ± 13.3%, 73.9 ± 3.0%, respectively, with each recovery level triplicated. The quantification result was summarized in Table 3.

The highest concentration of total alkaloid aglycone was found in S. villosum at a concentration less than 50 µg/g DW, or ~100 ug/g DW in terms of glycosylated forms, which was more than six times higher than the reported 1.62 \pm 0.02 mg/100 g DW in S. nigrum L. var. virginicum of Nigeria [20]. Assuming 90% moisture content in fresh vegetables, glycoalkaloid concentration was at maximum around 1 mg/100 g fresh weight (FW). In order to evaluate the potential risk of consumption, this level was compared with Solanum melongena or eggplant, a globally consumed vegetable well known to contain two principle glycoalkaloids, solasonine and solamargine, a same or similar composition as that found in AIVs in this study. An extensive study revealed a safe consumption of eggplants at glycoalkaloids levels ranging from 6.25 to 20.5 mg/100 g FW [25,26], which was significantly higher than the maximum glycoalkaloid level found in the AIVs in this research (p < 0.001). Given the low concentration of glycoalkaloids present in the leaves of the nightshades, results suggest that the consumption of these leafy AIVs are safe.

3.3. β -carotene and vitamin E analysis

β-Carotene and vitamin E in nightshade extracts were identified and quantified against standards using UV detection at 454 nm and 293 nm, respectively. Quantification results were summarized in Table 3. β -Carotene in all four species was higher than the average level of a wide scope of household fresh vegetables reported by USDA National Nutrient Database (Figure S1 and Table S1 of the supporting information). A total of 10 entries were identified to be a source of vitamin A (Table 3) based on retinol equivalent (RE) according to standards published by Codex Alimentarius [27,28]. The averaged β-carotene content across the five accessions of S. nigrum was $81.1\pm40.7\,\mu\text{g/g}\,\text{DW}.$ Except for S. nigrum PI 306400, all the other accessions contained higher level of β -carotene than that reported 4.66 \pm 0.02 mg/100 g DW or 46.6 \pm 0.2 μ g/g DW in a nutritive study of S. nigrum L. var. virginicum of Nigeria [20]. Up to 141.7 \pm 11.7 μ g/g DW, S. nigrum PI 312110 possessed the highest content in all accessions of S. nigrum and the other three species. In S. scabrum, SS 49 was found to contain the most abundant source of β -carotene with up to 96.0 ± 2.8 μ g/g DW. The mean level in all eight accessions of S. scabrum was $79.4 \pm 15.2 \,\mu$ g/g DW, nearly twice the level reported by Mibei at el. at 46 \pm 2.5 µg/g DW [29]. The β -carotene level in the single accession of S. americanum was $95.8 \pm 3.8 \mu g/g$ DW, less than a

fifth of the reported 52.1 \pm 3.6 mg/100 g DW in the literature [30]. In contrast, the β -carotene level of S. villosum was found to be high at 138.1 \pm 4.0 μ g/g DW and this is the report of β -carotene from this species.

Vitamin E content in all four species, except in a few entries, was among the top 25% when compared with the USDA National Nutrient Database (Fig. S1 and Table S1 of the supporting information). In the five accessions of S. *nigrum*, vitamin E content reached 150.3 \pm 59.9 µg/g DW on average. The accession S. *nigrum* PI 312110, was found not only to be high in β -carotene, but also the richest source of vitamin E among all accessions of the four studied species at 229.7 \pm 19.9 µg/g DW. In the eight entries of S. *scabrum*, the average level was 126.2 \pm 44.3 µg/g DW, with BG 16 having the highest concentration of vitamin. The content in the one accession of S. *americanum* was 145.5 \pm 5.2 µg/g, and in the single accession of S. villosum was 114.3 \pm 5.0 µg/g.

3.4. Total polyphenol content and antioxidant activity

The TPP level and antioxidant activity values were summarized in Table 3. The TPP level in the five accessions of S. *nigrum* averaged 10.07 \pm 2.40 mg GAE/g DW, and in the eight accessions of S. *scabrum* was 10.09 \pm 1.80 mg GAE/g DW. S. *scabrum* PI 643126 and Grif 14198 was reported in another study to contain free phenolic acids at a level of 12.83 \pm 0.41 mg GAE/g DW and 15.65 \pm 1.08 mg GAE/g DW, respectively, which was correspondingly 65% and 42.5% higher than the TPP level found in this study [31].

Antioxidant activity in the five accessions of S. *nigrum* was averaged to be 19.6 \pm 3.9 mg TE/g DW, and in the eight accessions of S. *scabrum* at 20.2 \pm 2.8 mg TE/g DW. In comparison, Jimenez-Aguilar using AAPH radical reported the antioxidant activity of S. *scabrum* PI 643126 and Grif 14198 to be 34.5 \pm 1.95 µmol TE/g FW and 36.17 \pm 1.47 µmol TE/g FW, respectively, or 50.15 \pm 2.83 mg/g DW and 51.35 \pm 2.09 mg/g DW in accordance [31]. Among samples in this study, there was a close correlation between TPP content and antioxidant capability with R² = 0.95 (n = 15), suggesting that either the TPP content or ABTS antioxidant activity may be used for quality control purpose.

4. Conclusion

Phytochemical screening by HPLC/ESI-MS showed that all four Solanum species contained abundant quantities of flavonoids and saponins. All 15 sources from the four Solanum species showed high levels of β -carotene, vitamin E and total polyphenol level and exhibited high antioxidant activity, suggesting that edible nightshades are nutrient rich and would be of health benefit. Importantly, as sub-Saharan Africans consume the leaves and not the fruits which have been long associated with anti-nutrient Solanaceous glycoalkaloids, no glycoalkaloids were detected except in two of the studied accessions, one from S. *nigrum* and another from S. *villosum*, where total solasodine aglycone was quantified after acid hydrolysis to be less than 50 ug/g DW, or 1 mg/100 g FW in terms of glycosylated forms. The potential consumption risk at this level is low as judged by comparison with that of globally consumed Solanum plant eggplant, whose glycoalkaloids were extensively studied. The glycoalkaloid content across all four species was found to be safe for consumption. The edible nightshades as part of a larger group of African indigenous vegetables were nutrient dense, possessing potential health benefits, and can be consumed without concern of glycoalkaloid related toxicity. Such results provide support for the promotion of their cultivation, marketing and consumption to provide available and accessible nutrient rich vegetables to improve household nutrition. The production of such vegetables could also provide income generating opportunities for those growers entering into the informal or formal markets.

Conflicts of interest

All contributing authors have no conflicts of interest to declare.

Acknowledgements

This research was supported by the Horticulture Innovation Lab with funding from the U.S. Agency for International Development (USAID EPA-A-00-09-00004), as part of the U.S. Government's global hunger and food security initiative, Feed the Future, for project titled "Improving nutrition with African indigenous vegetables" in eastern Africa. This study was made possible by the generous support of the American people through the United States Agency for International Development (USAID). The contents are the responsibility of the Horticulture Innovation Lab and do not necessarily reflect the views of USAID or the United States Government. This work is part of the senior authors dissertation research for food science. We thank Agilent Instruments, the New Jersey Agriculture Experiment Station (HATCH project 12131) and the World Vegetable Center. We also thank John Bowman, USAID-Washington, D.C., and Beth Mitcham, UC-Davis for their support and leadership in the Horticulture Innovation Lab.

Appendix A. Supplementary data

Supplementary data related to this article can be found at https://doi.org/10.1016/j.jfda.2017.10.005.

REFERENCES

- Keding G, Weinberger K, Swai I, Mndiga H. Diversity, traits and use of traditional vegetables in Tanzania. AVRDC-WorldVegetableCenter; 2007.
- [2] Weinberger K, Msuya JM. Indigenous vegetables in Tanzania: significance and prospects. AVRDC-WorldVegetableCenter; 2004.
- [3] Mal B. Neglected and underutilized crop genetic resources for sustainable agriculture. Indian J Plant Genet Resour 2007;20:1–14.

- [4] Yang R-Y, Ojiewo C. African nightshades and African eggplants: taxonomy, crop management, utilization and phytonutrients. Afr Nat Plant Prod 2013;2:137–65.
- [5] Milner SE, Brunton NP, Jones PW, O'Brien NM, Collins SG, Maguire AR. Bioactivities of glycoalkaloids and their aglycones from Solanum species. J Agric Food Chem 2011;59:3454–84.
- [6] Carle R. Investigations on the content of steroidal alkaloids and sapogenins within Solanum sect. Solanum (= sect. Morella)(Solanaceae). Plant Syst Evol 1981;138:61-71.
- [7] Zhen J, Villani TS, Guo Y, Qi Y, Chin K, Pan MH, et al. Phytochemistry, antioxidant capacity, total phenolic content and anti-inflammatory activity of *Hibiscus sabdariffa* leaves. Food Chem 2016;190:673–80.
- [8] Bai Y. Simultaneous quantification of four saponins, three alkaloids and three fatty acids in *Solanum nigrum Linn*. by HPLC-ELSD. J Med Plants Res 2012;6.
- [9] Do QD, Angkawijaya AE, Tran-Nguyen PL, Huynh LH, Soetaredjo FE, Ismadji S, et al. Effect of extraction solvent on total phenol content, total flavonoid content, and antioxidant activity of *Limnophila aromatica*. J Food Drug Anal 2014;22:296–302.
- [10] Crabbe PG, Fryer C. Modeling of acid hydrolysis of solasodine glycosides. Part III: combined glycoside hydrolysis and solasodiene formation - guidelines for commercial hydrolysis. AIChE J 1983;29:584–7.
- [11] Denardin CC, Hirsch GE, da Rocha RF, Vizzotto M, Henriques AT, Moreira JCF, et al. Antioxidant capacity and bioactive compounds of four Brazilian native fruits. J Food Drug Anal 2015;23:387–98.
- [12] Singleton V, Rossi JA. Colorimetry of total phenolics with phosphomolybdic-phosphotungstic acid reagents. Am J Enol Vitic 1965;16:144-58.
- [13] Kao F-J, Chiu Y-S, Chiang W-D. Effect of water cooking on antioxidant capacity of carotenoid-rich vegetables in Taiwan. J Food Drug Anal 2014;22:202–9.
- [14] Re R, Pellegrini N, Proteggente A, Pannala A, Yang M, Rice-Evans C. Antioxidant activity applying an improved ABTS radical cation decolorization assay. Free Radic Biol Med 1999;26:1231–7.
- [15] Nagulsamy P, Ponnusamy R, Thangaraj P. Evaluation of antioxidant, anti-inflammatory, and antiulcer properties of Vaccinium leschenaultii Wight: a therapeutic supplement. J Food Drug Anal 2015;23:376–86.
- [16] Matasyoh LG, Murigi HM, Matasyoh JC. Antimicrobial assay and phyto-chemical analysis of Solanum nigrum complex growing in Kenya. Afr J Microbiol Res 2014;8:3923–30.
- [17] Chen HJ, Inbaraj BS, Chen BH. Determination of phenolic acids and flavonoids in *Taraxacum formosanum* Kitam by liquid chromatography-tandem mass spectrometry coupled with a post-column derivatization technique. Int J Mol Sci 2012;13:260–85.
- [18] Guo S-B, Tian Y, Jian L-Y. Optimization of ethanol extraction process of Solanum nigrum Linn. and structural confirmation of its compounds. Asian J Chem 2014;26:4615.
- [19] Ding X, Zhu F, Yang Y, Li M. Purification, antitumor activity in vitro of steroidal glycoalkaloids from black nightshade (Solanum nigrum L.). Food Chem 2013;141:1181–6.
- [20] Akubugwo I, Obasi A, Ginika S. Nutritional potential of the leaves and seeds of black nightshade-Solanum nigrum L. var. virginicum from Afikpo-Nigeria. Pak J Nutr 2007;6:323–6.
- [21] Adesina S, Gbile Z. Steroidal constituents of Solanum scabrum subsp. nigericum. Fitoterapia 1984;55(6):362–3.
- [22] Mibei EK, Ojijo N, Karanja SM, Kinyua JK. Phytochemical and antioxidant analysis of methanolic extracts of four African indigenous leafy vegetables. Ann Food Sci Technol 2012;13:37–42.
- [23] Jared JJ, Murungi LK, Wesonga J, Torto B. Steroidal glycoalkaloids: chemical defence of edible African

nightshades against the tomato red spider mite, Tetranychus evansi (Acari: Tetranychidae). Pest Manag Sci 2016;72:828–36.

- [24] Vidal Aldana M, Noguiera Lima C. Isolation and characterization of a glycoside from fluid extracts of Solanum americanum Mill. Afinidad 1999;56:393–6.
- [25] Bajaj K, Kaur G, Chadha M. Glycoalkaloid content and other chemical constituents of the fruits of some eggplant (Solanum melongena. L.) varieties. J Plant Foods 1979;3(3):163–8.
- [26] Jones PG, Fenwick GR. The glycoalkaloid content of some edible solanaceous fruits and potato products. J Sci Food Agric 1981;32:419–21.
- [27] Food and Agriculture Organization of the United Nations WHO. Guidelines for use of nutrition and health claims (CAC/ GL 23–1997). Codex Alimentarius.

- [28] Food and Agriculture Organization of the United Nations WHO. Guidelines on nutrition labelling (CAC/GL 2–1985). Codex Alimentarius.
- [29] Mibei EK, Ojijo NKO, Karanja SM, Kinyua JK. Compositional attributes of the leaves of some indigenous African leafy vegetables commonly consumed in Kenya. Ann Food Sci Technol 2011;12:146–54.
- [30] Serrano J, Goni I, Saura-Calixto F. Determination of betacarotene and lutein available from green leafy vegetables by an in vitro digestion and colonic fermentation method. J Agric Food Chem 2005;53:2936–40.
- [31] Jimenez-Aguilar DM, Grusak MA. Evaluation of Minerals, phytochemical compounds and antioxidant activity of Mexican, central American, and African green leafy vegetables. Plant Foods Hum Nutr 2015;70:357–64.