

Species identification of a causative moray eel meat by SDS-PAGE

Follow this and additional works at: <https://www.jfda-online.com/journal>

Recommended Citation

Huang, Y.-C.; Chen, T.-Y.; Jeng, S.-S.; Chen, H.-M.; and Hwang, D.-F. (2010) "Species identification of a causative moray eel meat by SDS-PAGE," *Journal of Food and Drug Analysis*: Vol. 18 : Iss. 4 , Article 5.

Available at: <https://doi.org/10.38212/2224-6614.2238>

This Original Article is brought to you for free and open access by Journal of Food and Drug Analysis. It has been accepted for inclusion in Journal of Food and Drug Analysis by an authorized editor of Journal of Food and Drug Analysis.

Species Identification of a Causative Moray Eel Meat by SDS-PAGE

YUN-CHIEH HUANG¹, TAI-YUAN CHEN^{1,2}, SEN-SHYONG JENG¹,
HONG-MING CHEN³ AND DENG-FWU HWANG^{1*}

¹ Department of Food Science, National Taiwan Ocean University, Keelung, Taiwan, R.O.C.

² Institute of Biological Chemistry and Genomic Research Center, Academia Sinica, Taipei, Taiwan, R.O.C.

³ Department of Aquaculture, National Taiwan Ocean University, Keelung, Taiwan, R.O.C.

(Received: April 17, 2009; Accepted: March 30, 2010)

ABSTRACT

Attempts were made to identify the species of food poisoning moray eel. Besides the causative processed moray eel meat, eight other raw commercial moray eel meats including *Gymnothorax favagineus*, *G. fimbriatus*, *G. flavimarginatus*, *G. meleagris*, *G. pseudothyroideus*, *G. undulates*, *G. albimarginatus* and *G. javanicus* were heated at 100°C for 30 min and then assayed using the electrophoresis method of sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE). The SDS-PAGE patterns for eight processed commercial moray eel meats showed species-specific protein bands < 30 kD. The species of causative processed moray eel implicated in food poisoning case was identified as *G. javanicus* in comparison to the 2% SDS and 8 M urea extracted protein revealed by SDS-PAGE patterns.

Key words: moray eel, species identification, processed fish products, electrophoresis

INTRODUCTION

In April 2004, a food poisoning incident occurred due to ingesting cooked moray eel meat at Fugil fishing port, Taipei, Taiwan. A 47-year-old patient showed clinical symptoms of pricking sensation of lips, tongue and throat, vomiting, abdominal cramps, diarrhea, headache, dizziness, vertigo and paralysis. These symptoms disappeared gradually, and they were similar to those of ciguatera poisoning⁽¹⁾. Although ciguatera poisoning has been reported occasionally in Taiwan, the toxicity and the species of toxic fish samples implicated were rarely elucidated, because the causative residues had not been retained for study and the morphological characterization of the retained was difficult.

Moray eels are commercially important fish species in Taiwan, they are usually cut into fillets before selling. It is hard to identify the species by examining only the fillet morphology. These fishes are commonly found in the tropical and subtropical coral reef area around the world they are carnivorous and important edible species⁽²⁾. There are more than 20,000 cases of food poisoning

per year⁽³⁾. In 2001, one of the ciguateric toxins, ciguatoxin, has been isolated from moray eel *Gymnothorax javanicus*⁽²⁾. Ciguatoxin is produced by *Gambierdiscus toxicus*, which is usually attached on coral⁽⁴⁻⁶⁾. The toxin and related compounds are first accumulated by herbivorous fish and then transferred into carnivorous fish via food chain. When people consume these toxic herbivorous or carnivorous fish, they may be intoxicated⁽⁷⁾.

Since 2001, it has been shown that analysis of mitochondrial DNA (mtDNA) was successful in differentiating species of fish⁽⁸⁾. The cytochrome *b* (*cyt b*) gene has been used in species identification⁽⁹⁾ and in taxonomic phylogenetic studies⁽¹⁰⁾. The *cyt b* gene is considered one of the most useful genes for phylogenetic work and is probably the best-known mitochondrial gene with respect to structure and function of protein products^(11,12). Meanwhile, several electrophoretic methods have been employed to differentiate species of seafood or seafood products⁽¹³⁾. Other feasible methods have been subsequently applied in identifying fish species, including sodium dodecyl sulfate polyacrylamide gel electrophoresis (SDS-PAGE)⁽¹⁴⁻¹⁶⁾, urea-isoelectric focusing (urea-IEF)⁽¹⁷⁾, two-dimensional (2D) gel electrophoresis^(18,19), high performance liquid chromatography (HPLC)⁽²⁰⁾, immunoassay⁽²¹⁾ and

* Author for correspondence. Tel: +886-2-2462-2192 ext. 5103; Fax: +886-2-2462-6602; E-mail: dfhwang@mail.ntou.edu.tw

capillary electrophoresis (CE)⁽²²⁾.

The aims of this study were to establish basic data of SDS-PAGE patterns and to quantify the amount of SDS-extracted and urea-extracted proteins in eight commercial moray eel species in Taiwan. The species of causative processed moray eel fillet might then be identified compared to these commercial moray eel species.

MATERIALS AND METHODS

I. Fish Samples

Samples of eight moray eel species, *Gymnothorax favagineus* (GFA), *G. fimbriatus* (GFI), *G. flavimarginatus* (GFL), *G. meleagris* (GME), *G. pseudothyrsoides* (GPS), *G. undulates* (GUN), *G. albimarginatus* (GAL) and *G. javanicus* (GJA) were purchased from seafood markets in Keelung, Taiwan. Each species was represented by at least three specimens. The body weight and body length of tested moray eel samples are as follows: GFA, 541-1,313 g, 72-90 cm; GFI, 1,169-1,359 g, 87-90 cm; GFL, 938-2,617 g, 79-105 cm; GME, 1,054-1,363 g, 72-81 cm; GPS, 541-823 g, 67-74 cm; GUN, 1,022-1,303 g, 79-89 cm; GAL, 173-980 g, 49-103 cm; GJA, 554-3,350 g, 71-107 cm. Among them, the meat of each moray eel was collected from whole specimen and homogenized. A half of mixed meats were repeatedly sampled and rehomogenized. Finally 10 g of mixed meats was directly taken into centrifuge tubes and cooked at 100°C for 15 and 30 min, and 121°C for 15 min. After cooling, these heated samples and the causative samples (CS) of cooked moray eel fillet (250 g) provided by the victim were stored at -20°C until use.

II. Extraction Procedures

Two reagents (2% SDS and 8 M urea) were used. All extracts were supplemented with 0.1 mM phenylmethylsulfonyl fluoride (PMSF) and 0.01% (w/v) NaN₃ to inhibit proteases and microbial growth. The extracts of fish proteins were prepared according to the procedures described by Eitenne *et al.*⁽²³⁾ All operations were performed at room temperature.

For the SDS extracts, 1.0 g of heated samples were homogenized with 5 mL of the extraction solution using a Polytron-Aggregate (setting 4.0, Kinematica Littau/Luzern, Switzerland) at low speed for 1 min. After homogenizing, the mixture was boiled for 2 min in a water bath at 100°C, and then homogenized again. The homogenates were kept at room temperature for 30 min and then centrifuged at 10,000 ×g for 15 min at 10°C using Hima CF 15D2 with rotor RT15A6 (Hitachi, Japan). The supernatants were collected and stored at room temperature (maximum one week) before analyzed by electrophoresis.

For the urea extracts, the same procedure for

homogenizing was followed. After homogenizing, the homogenates were kept at room temperature for 30 min, and then centrifuged as described before. The supernatants were collected, and then stored as SDS extracts.

III. Determination of Protein Concentration

The protein concentration was determined by bicinchoninic acid method⁽²⁴⁾. The protein extracts were adjusted with sample buffer to 2 mg/mL for electrophoresis.

IV. SDS-PAGE

SDS-PAGE was performed according to the modified procedure of Laemmli⁽²⁵⁾ and O'Farrell⁽²⁶⁾, using a Mini-Protean unit (Bio-Rad, Richmond, CA, USA). Slab gels consisted of a separating gel (15.0%), which was polymerized for 1 hr, and a stacking gel (4.0%), which was poured for 30 min before sample application. The 10 µL of samples (20 µg of extracted protein) were applied in the wells of the gel. Electrophoresis was carried out at constant voltage of 150 volts when the tracking dye reached the separating gel. Electrophoresis was completed when the dye front reached the bottom of the gel.

V. Gel Stain and Destain

Gels were stained either with Coomassie blue reagent, 0.1% (w/v) Coomassie blue R-250 in 40% methanol and 10% acetic acid. After staining, gels were destained in 10% methanol and 10% acetic acid. Molecular weights were determined by comparing relative mobilities of protein bands to standard proteins⁽²⁷⁾. Protein standards were obtained from Bio-Rad (Broad range kit: myosin 211 kD, β-galactosidase 121 kD, bovine serum albumin 100 kD, ovalbumin 54.4 kD, carbonic anhydrase 38.7 kD, soybean trypsin inhibitor 29.8 kD, lysozyme 20.0 kD, aprotinin 7.3 kD).

VI. Image Analysis

The gels were scanned by Gel Dox[®] (Bio-Rad) and the acquired images were analyzed with software Quantity One[®] (Bio-Rad). Images of protein profiles were stored in computer and molecular weights were estimated by comparing R_f values on the gel with those of the protein standards. The relative amount of each protein band in SDS-PAGE patterns was the optical density of each protein band to total optical densities of all protein bands in the same lane.

RESULTS AND DISCUSSION

The amounts of SDS-extracted proteins and urea-extracted proteins from moray eel meat were shown

in Table 1. There were significant differences between the raw fish and the ones heated at 100°C for 15 min, 100°C for 30 min and 121°C for 15 min in SDS-extracted proteins, showing the decrease of 7.4%, 21.3% and 35.0%, respectively. The results of urea-extracted proteins showed greater decrease in protein amount with 19.5%, 36.9% and 47.3%, respectively. Thus the increase in heating temperature and time both reduced the extractable proteins from raw moray eel meat. It also showed that the protein concentration of SDS-extracted was higher than that of urea-extracted. It indicated that 2% SDS was more suitable than 8 M urea to extract denatured protein from processed eels. Figure 1 shows the

SDS-PAGE patterns of SDS-extracted proteins (Figure 1A) and urea-extracted proteins (Figure 1B) from *G. javanicus* stained with Coomassie blue after different processed conditions. At the heat-processing temperature of 100°C for 15 min and 30 min, the protein bands of SDS-extracted and urea-extracted proteins were clear and remained differentiable. However, the processed moray eel meats heated at 121°C for 15 min were degraded seriously and could not be used in species identification (Figure 1).

Judging from the high molecular weight region (> 30.0 kD) of two extracted proteins stained with Coomassie blue, there were no species-specific protein bands among





Figure 1. SDS-PAGE patterns of moray eel protein extracted with SDS (A) and urea (B) from *G. javanicus* meat with different heating processes. A: 100°C, 15 min; B: 100°C, 30 min; C: 121°C, 15 min. M: protein standards; R: raw-meat of *G. javanicus*.

Table 1. The protein amounts (n = 3) of SDS and urea extracts from raw and processed moray eel meat

Sample	SDS extract (mg)				Urea extract (mg)			
	Raw	Processed meat			Raw	Processed meat		
		100°C, 15 min	100°C, 30 min	121°C, 15 min		100°C, 15 min	100°C, 30 min	121°C, 15 min
GFA	117.5 ± 0.1	107.7 ± 0.1	88.0 ± 0.1	68.5 ± 0.1	111.5 ± 0.1	77.0 ± 0.1	63.5 ± 0.1	51.5 ± 0.1
GFI	119.0 ± 0.1	106.0 ± 0.1	84.5 ± 0.1	71.5 ± 0.1	104.0 ± 0.1	87.5 ± 0.1	61.5 ± 0.1	56.0 ± 0.1
GFL	109.5 ± 0.1	103.0 ± 0.1	90.5 ± 0.1	80.5 ± 0.1	104.0 ± 0.1	80.5 ± 0.1	68.0 ± 0.1	47.5 ± 0.1
GME	114.5 ± 0.1	106.5 ± 0.1	87.5 ± 0.1	66.5 ± 0.1	97.0 ± 0.1	66.5 ± 0.1	52.0 ± 0.1	52.5 ± 0.1
GPS	114.0 ± 0.1	105.0 ± 0.1	89.5 ± 0.1	73.0 ± 0.1	91.0 ± 0.1	85.5 ± 0.1	49.0 ± 0.1	50.0 ± 0.1
GUN	114.0 ± 0.1	103.0 ± 0.1	98.5 ± 0.1	76.0 ± 0.1	93.0 ± 0.1	87.0 ± 0.1	48.5 ± 0.1	55.0 ± 0.1
GAL	100.0 ± 0.1	95.5 ± 0.1	84.0 ± 0.1	65.5 ± 0.1	94.0 ± 0.1	74.5 ± 0.1	61.0 ± 0.1	49.0 ± 0.1
GJA	113.0 ± 0.1	107.7 ± 0.1	86.0 ± 0.1	83.5 ± 0.1	95.5 ± 0.1	74.5 ± 0.1	94.5 ± 0.1	53.0 ± 0.1
CS*			63.0 ± 0.1				49.5 ± 0.1	

*CS (causative sample) was the moray eel fillet provided by the victim.



MW kD	GFA	GFI	GFL	GME	GPS	GUN	GAL	GJA	CS
29.0							2.58 ± 0.07		
28.7			0.57 ± 0.04	0.84 ± 0.03		3.95 ± 0.07			
28.4					2.05 ± 0.04				
27.7			0.58 ± 0.03						
26.9				2.23 ± 0.03			1.01 ± 0.05		
26.7	2.36 ± 0.07								
26.2		0.67 ± 0.06							
25.9								4.80 ± 0.03	4.75 ± 0.02
25.7						4.60 ± 0.03			
25.4							2.55 ± 0.03		
25.1					4.60 ± 0.03				
23.0			0.73 ± 0.05	2.10 ± 0.07					
22.4							3.33 ± 0.08	4.45 ± 0.06	4.62 ± 0.05
22.1	4.10 ± 0.07	2.65 ± 0.07			4.01 ± 0.02	6.00 ± 0.02			
20.5			0.50 ± 0.03						
16.4				0.81 ± 0.02					
14.5						3.09 ± 0.03			
14.2		0.47 ± 0.04					0.81 ± 0.03	1.25 ± 0.04	0.82 ± 0.05
13.5					0.73 ± 0.02				
9.3					3.39 ± 0.05				
8.9	4.67 ± 0.03		0.83 ± 0.03						
8.5				5.20 ± 0.07					
8.3		7.01 ± 0.02							
8.0	1.50 ± 0.03		0.91 ± 0.04						
7.7								8.71 ± 0.04	4.95 ± 0.03
7.3						6.14 ± 0.08	3.72 ± 0.04		
6.1					2.15 ± 0.04				


Figure 2. SDS-PAGE patterns of SDS extracted protein from eight moray eel species (100°C, 30 min) stained with Coomassie blue (n = 3). M: protein standards, GFA: *Gymnothorax favagineus*, GFI: *G. fimbriatus*, GFL: *G. flavimarginatus*, GME: *G. meleagris*, GPS: *G. pseudothyrsoideus*, GUN: *G. undulatus*, GAL: *G. albimarginatus*, GJA: *G. javanicus*, and CS: causative sample. Data shown as Mean ± STD.

these moray eel species (Figure 2). In addition, the amounts of high molecular weight region (> 30.0 kD) were too large to be differentiated between moray eel species. The bands at the high molecular weight region were apparently not useful in identifying moray eel species.

All tested moray eel species could be differentiated using the SDS-PAGE patterns of SDS-extracted proteins in the low molecular weight region (< 30.0 kD) (Figure 2). The causative moray eel meat established the same

SDS-PAGE patterns of the SDS-extracted proteins of *G. javanicus*. Therefore, SDS-PAGE patterns of the SDS-extracted protein stained with Coomassie blue would be useful for moray eel species identification.

The low molecular weight region (< 30.0 kD) of the SDS-PAGE patterns of urea-extracted proteins showed species-specific protein patterns (Figure 3). The percentage compositions of urea-extracted proteins of the low molecular weight (< 30.0 kD) are shown in Figure 3.



MW kD	GFA	GFI	GFL	GME	GPS	GUN	GAL	GJA	CS
27.4		0.65 ± 0.03							
27.1					2.74 ± 0.04	0.41 ± 0.02	0.50 ± 0.04		
26.1			1.43 ± 0.06					0.79 ± 0.11	1.42 ± 0.11
25.5	1.92 ± 0.02								
23.1		5.93 ± 0.05					4.50 ± 0.03		
22.9					5.73 ± 0.04	3.38 ± 0.08			
22.2			4.27 ± 0.07	5.53 ± 0.07				4.77 ± 0.04	5.81 ± 0.15
21.9	9.97 ± 0.06								
16.7		3.20 ± 0.08					0.57 ± 0.06		
16.3					2.51 ± 0.07				
15.7						0.53 ± 0.05			
15.4			1.84 ± 0.01	3.67 ± 0.10					
13.3	3.46 ± 0.02								
10.4					5.65 ± 0.04				
8.7	5.34 ± 0.01		5.33 ± 0.05						
8.3		7.97 ± 0.03			2.21 ± 0.03		4.66 ± 0.07	7.75 ± 0.12	5.77 ± 0.08
8.1						8.16 ± 0.09			
7.8	3.50 ± 0.05		4.03 ± 0.09	7.19 ± 0.12					

Figure 3. SDS-PAGE patterns of urea extracted proteins from eight moray eel species (100°C, 30 min) stained with Coomassie blue (n = 3). The symbols refer to moray eel species and the footnote detailed in Figure 2.

The moray eel species could be identified by the pattern of the low molecular weight region in the urea-extracted proteins. The causative moray eel meat showed similar SDS-PAGE pattern of the urea-extracted proteins to that of *G. javanicus*. Hence, the species of the causative moray eel was identified as *G. javanicus*.

The low molecular weight region of the SDS-PAGE patterns of solvent extracted proteins which shows species-specific protein patterns were similar to other reports^(4,15,17). In this study, the individual mobility variation between fish of the same species was not measured. Protein band density appeared somewhat variable and this may have a slight effect on our ability to discriminate moray eel species.

Moray eels have relatively higher price in Taiwan. Therefore, some fraudulent retailers use cheaper fish to substitute for moray eels. Because moray eel products are usually heavily processed, the proteins in the processed moray eel products have usually degraded and almost all proteins can be denatured and damaged, such as by heating at 121°C for 15 min. Then, the protein analysis methods for fish species identification are inappropriate. In this situation, polymerase chain reaction (PCR) amplification and restriction enzyme analysis of the cytochrome *b* gene may be applied for identification of fish species^(11,28).

CONCLUSIONS

Species identification of 8 tested moray eel species can be achieved by examining the lower molecular weight protein bands following SDS-PAGE of 2 % SDS or 8 M urea extractants along with the densities of the characteristic protein bands using staining. The below 30.0 kD protein bands in SDS-PAGE pattern were more useful than those of higher molecular weight proteins (> 30.0 kD). Moreover, the causative heated moray eel fillet implicated in the food poisoning was identified as *G. javanicus*.

ACKNOWLEDGMENTS

Moray eels were identified by Dr. Kang-Tsao Shao, Institute of Cellular and Organismic Biology, Academia Sinica, Taipei, Taiwan, R.O.C. and by co-author Dr. Hong-Ming Chen using morphological analysis. This study was partly supported by National Science Council (NSC-95-2313-B-019-024-MY3) and Center for Marine Bioscience and Biotechnology, National Taiwan Ocean University (9552900 1A4).

REFERENCES

1. Hashimoto, Y. 1979. Marine Toxins and Other Bioactive Marine Metabolites. pp. 91-158. Japan Scientific Societies, Tokyo, Japan.
2. Lewis, R. J. 2001. The changing face of ciguatera. *Toxicon* 39: 97-106.
3. Ito, E., Toyota, F. S., Toshimori, K., Fuwa, H., Kazuo, T., Satake, K. and Sasaki, M. 2003. Pathological effects on mice by gambierol, possibly one of the ciguatera toxins. *Toxicon* 42: 733-740.
4. Bagnis, R., Chanteau, S., Chungue, E., Hurtel, J. M., Yasumoto, T. and Inoue, A. 1980. Origins of ciguatera fish poisoning: a new dinoflagellate, *Gambierdiscus toxicus* Adachi and Fukuyo, definitively involved as a causal agent. *Toxicon* 18: 199-208.
5. Holmes, M. J., Lewis, R. J., Poli, M. A. and Gillespie, N. C. 1991. Stain dependant production of ciguatoxin precursors (gambiertoxin) by *Gambierdiscus toxicus* (Dinophyceae) in culture. *Toxicon* 29: 761-775.
6. Poli, M. A., Lewis, R. J., Dickey, R. W., Musser, S. M., Buckner, C. A. and Carpenter, L. G. 1997. Identification of Caribbean ciguatoxins as the cause of an outbreak of fish poisoning among U.S. soldiers in Haiti. *Toxicon* 35: 733-741.
7. Colman, J. R., Dechranoui, M. Y. B., Dickey, R. W. and Ramsdell, J. S. 2002. Characterization of the developmental toxicity of Caribbean ciguatoxins in finfish embryos. *Toxicon* 44: 59-66.
8. Quinteiro, J., Vidal, R., Izquierdo, V., Sotelo, C. G., Chapela, M. J. and Perez-Martin, R. I. 2001. Identification of hake species (*Merluccius geneus*) using sequencing and PCR-RFLP analysis of mitochondrial DNA control region sequences. *J. Agric. Food Chem.* 49: 5108-5114.
9. Cook, C. E., Wang, Y. and Sensabaugh, G. 1999. A mitochondrial control region and cytochrome *b* phylogeny of sika deer (*Cervus nippon*) and report of tandem repeats in the control region. *Mol. Phylogenet. Evol.* 12: 47-56.
10. Kuwagyjam, R. and Ozawa, T. 2000. Phylogenetic relationships among European red deer, wapiti, and sika deer inferred from mitochondrial DNA sequences. *Mol. Phylogenet. Evol.* 15: 115-123.
11. Farias, I. P., Orti, G., Sampaol, I., Schneider, H. and Meyer, A. 2001. The cytochrome *b* gene as a phylogenetic marker: The limits of resolution for analyzing relationship among cichlid fish. *J. Mol. Evol.* 53: 89-103.
12. Tsai, W. L., Chen, H. M., Hsieh, C. H., Lin, W. F. and Hwang, D. F. 2009. A potential methodology for differentiation of ciguatoxin-carrying species of moray eel. *Food Control* 20: 575-579.
13. Hollingworth, T. and Wekell, M. M. 1990. Official Methods of Analysis of the Association of Official Analytical Chemist. 15th ed. pp. 885-886. Association of Official Analytical Chemists. Arlington, Virginia, U.S.A.
14. Pineiro, C., Barros-Velazquez, J. R., Perez-Martin, I., Martinez, I., Jacobsen, T., Rehbein, H., Kundiger, R., Mendes, R., Etienne, M., Jerome, M., Craig, A.,

- Mackie, I. M. and Jessen, F. 1999. Development of a sodium dodecyl sulfate- polyacrylamide gel electrophoresis reference method for the analysis and identification of fish species in raw and heat-processed samples: A collaborative study. *Electrophoresis* 20: 1425-1432.
15. Chen, T. Y. and Hwang, D. F. 2002. Electrophoretic identification of muscle proteins in 7 puffer species. *J. Food Sci.* 67: 936-942.
16. Chen, T. Y., Hsieh, Y. W., Tsai, Y. H., Shiau, C. Y. and Hwang, D. F. 2002. Identification of species and measurement of tetrodotoxin in dried dressed fillets of the puffer fish, *Lagocephalus lunaris*. *J. Food Prot.* 65: 1670-1673.
17. Chen, T. Y., Shiau, C. Y., Noguchi, T., Wei, C. I. and Hwang, D. F. 2003. Identification of puffer fish species by native isoelectric focusing technique. *Food Chem.* 83: 475-479.
18. Martinez, I. and Friis, T. J. 2004. Application of proteome analysis to seafood authentication. *Proteomics* 4: 347-354.
19. Chen, T. Y., Shiau, C. Y., Wei, C. I. and Hwang, D. F. 2004. Preliminary study on puffer fish proteome-species identification on puffer fish by two-dimensional electrophoresis. *J. Agric. Food Chem.* 52: 2236-2241.
20. Wang, Q., Ruan, X., Jin, Z. H., Yan, Q. C. and Tu, S. 2005. Identification of *Rhodiola* species by using RP-HPLC. *J. Zhejiang Univ. Sci. B* 6: 477-482.
21. Lopata, A. L., Luijck, T., Fenemore, B., Sweijid, N. A. and Cook, P. A. 2002. Development of a monoclonal antibody detection assay for species-specific identification of abalone. *Mar. Biotechnol.* 4: 454-462.
22. Vallejo-Cordoba, B. and Cota-Rivas, M. 1998. Meat species identification by linear discriminant analysis of capillary electrophoresis protein profiles. *J. Capillary Electrophor.* 5: 171-175.
23. Eitenne, M., Jerome, M., Fleurence, J., Rehbein, H., Kundiger, R., Yman, I. M., Ferm, M., Craig, A., Mackie, I., Jessen, F., Smelt, A. and Lutten, J. 1999. A standardized method of identification of raw and heat-processed fish by urea isoelectric focusing: a collaborative study. *Electrophoresis* 20: 1923-1933.
24. Smith, P. K., Krohn, R. I., Hermanson, G. T., Mallia, A. K., Gartner, F. H., Provenzano, M. D., Fujimoto, E. K., Geoke, N. M., Olson, B. J. and Klenk, D. C. 1985. Measurement of protein using bicinchoninic acid. *Anal. Biochem.* 150: 76-85.
25. Laemmli, U. K. 1970. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 227: 680-685.
26. O'Farrell, P. H. 1975. High resolution two-dimensional electrophoresis of proteins. *J. Biol. Chem.* 250: 4007-4021.
27. Weber, K., Pringle, J. R. and Osborn, M. 1972. Measurement of molecular weights by electrophoresis on SDS-acrylamide gel. *Meth. Enzymol.* 26: 3-27.
28. Hsieh, Y. W. and Hwang, D. F. 2004. Molecular phylogenetic relationships of puffer fish inferred from partial sequence of cytochrome *b* gene and restriction fragment length polymorphism analysis. *J. Agric. Food Chem.* 52: 4159-4156.