

Inhibition of benzo(a)pyrene hydroxylation by lignans isolated from *Justicia procumbens*

Follow this and additional works at: <https://www.jfda-online.com/journal>

Recommended Citation

Ueng, Y.-F.; Chen, C.-C.; and Chen, C.-F. (2000) "Inhibition of benzo(a)pyrene hydroxylation by lignans isolated from *Justicia procumbens*," *Journal of Food and Drug Analysis*: Vol. 8 : Iss. 4 , Article 9.
Available at: <https://doi.org/10.38212/2224-6614.2822>

This Original Article is brought to you for free and open access by Journal of Food and Drug Analysis. It has been accepted for inclusion in Journal of Food and Drug Analysis by an authorized editor of Journal of Food and Drug Analysis.

Inhibition of Benzo(a)pyrene Hydroxylation by Lignans Isolated from *Justicia procumbens*

YUNE-FANG UENG*, CHIEN-CHIH CHEN AND CHIEH-FU CHEN

National Research Institute of Chinese Medicine, Taipei, Taiwan, R. O. C.

(Received: June 26, 2000; Accepted: September 6, 2000)

ABSTRACT

Five lignans, neojustin A (neo A), neojustin B (neo B), justicidin A (jus A), justicidin B (jus B), and chinensinaphthol methyl ether (CME) were isolated from the ethanol extract of *Justicia procumbens* (*J. p.*), which has been used as a herbal remedy in traditional Chinese medicine. The *in vitro* effects of lignans on rat hepatic cytochrome P450-catalyzed oxidations were studied. Addition of 10 μM lignans caused 21% to 57% decreases of microsomal benzo(a)pyrene hydroxylation (AHH) activity. Among these lignans, neo B had the strongest inhibitory effect on AHH activity with an IC_{50} of $6 \pm 1 \mu\text{M}$. Lignans at 10 μM decreased 7-ethoxyresorufin O-deethylation activities by 20% to 48%. Neo B at 10 μM caused a 70% decrease of 7-methoxyresorufin O-demethylation activity whereas this oxidation activity was relatively less affected by other lignans. Lignans (10 μM) also decreased testosterone 6 β -hydroxylation activity by 19% to 74% and neo B had the least inhibitory effect on this activity. Kinetic analysis of AHH activity revealed that neo B was a mixed type inhibitor of competitive and noncompetitive characteristics with K_i and K_i' of 2 μM and 11 μM , respectively. With NADPH as the variable cofactor, neo B showed a uncompetitive type of inhibition of AHH activity. These *in vitro* results suggested that lignans from *J. p.* inhibited monooxygenase activities differentially and neo B might have a role in diminishing the oxidative activation of benzo(a)pyrene.

Key words: *Justicia procumbens*, lignan, benzo(a)pyrene hydroxylation

INTRODUCTION

The cytochrome P450 (CYP)-dependent monooxygenase system consists of CYP, NADPH-CYP reductase, and phospholipids. This monooxygenase plays a major role in the detoxification and bioactivation of xenobiotics including drugs, environmental pollutants, and chemical carcinogens (1,2). Benzo(a)pyrene (B(a)P), a polycyclic aromatic hydrocarbon (PAH), is released into the atmosphere from incomplete combustion processes. Humans ingest B(a)P and other PAHs from charbroiled meat, cigarette smoke, and automobile emissions (3). CYP-mediated metabolic activation of B(a)P is necessary for its conversion to form the reactive metabolite, B(a)P 7,8-diol-9,10-epoxide, which binds to DNA and initiates tumor formation (4). The importance of the metabolic activation of B(a)P has led to the suggestion that inhibitors of B(a)P hydroxylation (AHH) activity could diminish the mutagenicity and carcinogenicity of B(a)P (5,6). Therefore, we examined the inhibition of AHH activity by traditional Chinese medicine.

Justicia procumbens (*J. p.*) has been used as a traditional herbal remedy for the treatment of fever and pain from pharyngolaryngeal swelling. Recently, it has been used as a constituent of a herbal tea in Taiwan. Arylnaphthalene lignans have been reported as the major active components in *J. p.* (7,8). These lignans also showed antiviral and antitumor activities (9). Five lignans (Scheme 1), neojustin A (neo A), neojustin B (neo B), justicidin A (jus A), justicidin B (jus B), and chinensinaphthol methyl ether (CME) have been iso-

lated and purified from the ethanol extract of *J. p.* by monitoring the anti-platelet activity (10). To investigate the beneficial effects of these lignans against B(a)P, the effects of lignans on microsomal AHH and related monooxygenase activities were studied *in vitro*.

MATERIALS AND METHODS


I. Lignans and Chemicals

Lignans were isolated from air-dried *J. p.* by the method described previously (10). The purity of these lignans were > 95% as analyzed by NMR and HPLC analyses. NADP, NADPH, glucose-6-phosphate, glucose-6-phosphate dehydrogenase, cytochrome *c*, B(a)P, 7-ethoxyresorufin, 7-methoxyresorufin, testosterone and 6 β -hydroxytestosterone were purchased from Sigma Chemical Co. (St. Louis, MO, U.S.A.). Methanol, acetone, *n*-hexane, and dichloromethane were purchased from Merck Chemical Co. (Darmstadt, Germany).

II. Animal Treatments

Male Sprague Dawley rats (four weeks, weighing 200 ~ 250 g) were purchased from the Animal Center of National Yang-Ming University, Taipei, Taiwan. Microsomes were isolated from rat livers by differential centrifugation (11). Washed microsomal pellets were covered with 0.1 M potassium phosphate buffer, pH 7.4 and stored at -75°C . Monooxygenase activities were determined within two weeks.

* Author for correspondence. Tel: 02-28201999 ext. 6351; Fax: 02-28264266; E-mail: ueng@cma23.nricm.edu.tw



Scheme 1. Structures of arylnaphthalene lignans isolated from *Justicia procumbens*⁽¹⁰⁾.

III. Monooxygenase Assays

The CYP contents were determined by CO-difference spectral analysis as described by Omura and Sato⁽¹²⁾. NADPH-CYP reductase activity was determined following the method of Phillips and Langdon⁽¹³⁾ using cytochrome *c* as a substrate. In the AHH assay, formation of 3-hydroxybenzo(a)pyrene was detected by fluorometry⁽¹⁴⁾. Activities of 7-ethoxyresorufin O-deethylation (EROD) and 7-methoxyresorufin O-demethylation (MROD) were determined by measuring fluorescence of resorufin⁽¹⁵⁾. For testosterone 6 β -hydroxylation (TESH), the formation of 6 β -hydroxytestosterone was analyzed by HPLC using a C18 column (100 Å, 4.6 \times 250 mm, Varian Analytical Instrument, CA, U.S.A.) and detected by A_{254nm}⁽¹⁶⁾. Microsomal protein concentration was determined by the method described by Lowry *et al.*⁽¹⁷⁾.

IV. Data Analysis

The IC₅₀ value of neo B for AHH activity was calculated by graph fitting (Grafit, Erithacus Software Ltd., Staines, UK). Kinetic analysis of AHH activity was done following the Michaelis-Menten kinetic property. Values of velocity (*v*) at various substrate concentrations (*s*) were fitted by nonlinear least-squares regression without weight due to the equation, $v = V_m(s/(K_m + s))$ (Sigma plot, Jandel Scientific, San Rafael, CA). The mixed type inhibitor constants for enzyme (*K_i*) and enzyme-substrate complex (*K_i'*) were obtained by linear regression of the plots of *I* vs 1/*V_m*(*i*) and *I* vs *K_m*(*i*)/*V_m*(*i*), where *I* was the concentrations of neo B and *V_m*(*i*) and *K_m*(*i*) were *V_m* and *K_m* in the presence of neo B,

respectively. Values of velocity at various NADPH concentrations were fitted by nonlinear least-squares regression without weight due to the equation, consistent with uncompetitive inhibition according to the Michaelis-Menten equation: $v = V_m \cdot S / \{K_m + S \cdot [1 + (I/K_i)]\}$. Estimates of variances (denoted by \pm) are presented from analysis of individual sets of data.

RESULTS AND DISCUSSION

To study the effects of lignans on CYP-catalyzed oxidations, monooxygenase activities were determined in the presence of increasing concentrations of lignans (Fig. 1). The lignans at 10 μ M inhibited rat microsomal AHH activity by 21% to 57% (Fig. 1A). Among these lignans, neo A and neo B showed greater inhibitory effects on AHH activity and neo B had the most potent inhibitory effect on AHH activity with an IC₅₀ of 6 \pm 1 μ M. Many studies have been done on rat hepatic CYP enzymes and have suggested that CYPs including CYP1A, CYP2C, and CYP3A were involved in the oxidation of B(a)P⁽¹⁸⁾. CYP1A has the highest AHH activity and CYP3A is the most abundant CYP in mammalian livers^(18,19,20). CYP1A1 and CYP1A2 are two major members of the CYP1A subfamily. CYP1A2 is the main hepatic CYP1A member and CYP1A1 is essentially an extrahepatic CYP1A in rodents and humans⁽²⁰⁾. EROD and MROD were preferentially catalyzed by CYP1A1 and CYP1A2, respectively^(21,22). In control rats, EROD was mainly catalyzed by CYP2C and CYP2B⁽²²⁾. Our data showed that 10 μ M lignans caused 20% to 48% decreases of EROD activity (Fig. 1B). The inhibitory effects of lignans were jus B > CME \simeq neo B > neo A \simeq jus A. The presence of neo B inhibited MROD




Figure 1. Inhibitory effects of aryl-naphthalene lignans on monooxygenase activities of rat liver microsomes. Increasing concentrations of lignans (○, neo A; ●, neo B; △, jus A; ▲, jus B; □, CME) were added to the NADPH-supported microsomal reaction mixtures and monooxygenase activities were determined as described in the section of Materials and Methods.

activity in a concentration-dependent manner, whereas there were relatively less inhibitory effects from the other lignans (Fig. 1C). Neo B at 10 μM caused a 70% decrease of MROD activity. TESH was selectively catalyzed by CYP3A⁽²³⁾. Lignans (10 μM) decreased TESH by 19% to 74%. The inhibitory effects of lignans were jus A > jus B \approx neo A \approx CME > neo B (Fig. 1D). Neo B had the least inhibitory effect on TESH activity. Our results suggest that these lignans could affect CYP activities with differential potency and neo B had stronger inhibitory effects on both AHH and MROD activities. The relationship between structures and the inhibitory effects of lignans is not clear. However, neo B showed the strongest inhibition on MROD activity, but the least inhibition on TESH activity. Jus A and jus B had a stronger inhibition effect on TESH activity, but less inhibition on AHH and MROD activities. These results suggest that the relative position of the C ring and the keto group in the γ -lactone moiety and the presence of 4-, 6- and 7-methoxy groups might play roles in the determination of inhibition selectivity of CYP1A and CYP3A.

To understand the inhibitory property of neo B on AHH

Table 1. Kinetic analyses of AHH activity in the presence of neo B

Neo B, μM	V_m	K_m	r^2 *
0	415 \pm 13	11 \pm 4	0.95
1	316 \pm 13	9 \pm 2	0.99
5	251 \pm 16	12 \pm 4	0.97
10	200 \pm 19	12 \pm 6	0.94

Hepatic AHH activity and kinetic parameters were determined as described in the section of Materials and Methods. * r^2 represents coefficient of the nonlinear regression determination.

activity, kinetic analysis of AHH activity in the presence of various concentrations of neo B were performed. Fig. 2A shows the velocity (v) vs substrate concentration (s) plot. Nonlinear regression analysis generated the V_m and K_m values in the presence of increasing concentrations of neo B (Table 1). In the presence of 10 μM neo B, the V_m value was decreased to 48% of the value in the absence of neo B (Table 1). In the presence of 1, 5, and 10 μM neo B, the K_m values were 9 \pm 2, 12 \pm 4, and 12 \pm 6 μM , respectively (Table 1). Kinetic analysis of the Lineweaver-Burk plots suggested that neo B was a mixed type of competitive and noncompetitive




Figure 2. Steady-state kinetics of AHH activity and the effects of neo B. Plots of v , expressed in pmol/min/mg protein, vs S , benzo(a)pyrene concentration (A) and Lineweaver-Burk plots (B) at increasing concentrations of neo B (\circ , 0 μM ; \bullet , 1 μM ; \blacktriangle , 5 μM ; \blacksquare , 10 μM) are presented, with lines drawing according to the fitting results using Sigma-Plot program as described in the section of Materials and Methods. Plots of $1/v$ vs neo B concentration (C) at various concentrations of benzo(a)pyrene (\circ , 10 μM ; \bullet , 30 μM ; \blacktriangle , 100 μM).




Figure 3. Kinetic analysis of the inhibition of AHH by neo B with respect to the cofactor NADPH. Plots of v , expressed in pmol/min/mg protein, vs NADPH concentrations (A) and Lineweaver-Burk plots (B) at increasing concentrations of neo B (\circ , 0 μM ; \bullet , 5 μM ; \blacktriangle , 10 μM).

inhibitors (Fig. 2B). The inhibitor constants of neo B, K_i and K_{i1} for the enzyme and enzyme-substrate complex were obtained by linear regression (the inserted figures in Fig. 2A and B; Fig. 2C). The K_{i1} value was larger than the K_i value (Table 2). This result also suggested that neo B was a mixed type inhibitor. This mixed type inhibitor suggests that neo B may bind to both CYP and CYP-substrate complex and interfere with the substrate binding and product formation. In supporting CYP-catalyzed oxidations, NADPH serves as an electron donor to NADPH-CYP reductase. Kinetic analysis of AHH activity was performed in conditions with varying concentrations of NADPH. With NADPH as the variable substrate, neo B showed an uncompetitive type of inhibition in AHH (Fig. 3). The inhibition constant K_i calculated by nonlinear regression was $7 \pm 1 \mu\text{M}$ (Table 2).

Several possibilities may contribute to the *in vitro* inhibition of neo B. First, neo B may have caused protein

Table 2. The inhibitor constants of neo B for AHH activity with respect to varying concentrations of benzo(a)pyrene and NADPH

Varying factor	Constant	μM	r^2 *
Benzo(a)pyrene	K_i	2	0.99
	K_{i1}	11	0.96
NADPH	K_i	7 ± 1	0.96

Inhibitor constants were determined as described in the section of Materials and Methods. * r^2 represents the coefficient of regression determination.

destruction and resulted in the inhibition of catalytic activity. To reveal this possibility, microsomes were incubated with or without 10 μM neo B for 10 min at room temperature and the CO-difference spectra were then determined. The microsomal CO-difference spectra were not affected by the treatment of lignans. This result suggested that lignans did not cause the destruction of the active heme moiety of CYP. Second,

neoB may interfere with the electron transfer from NADPH through NADPH-CYP reductase by reducing NADPH-CYP reductase activity. To examine this possibility, NADPH-CYP reductase activity toward cytochrome *c* was determined. However, the addition of neo B in the reaction mixture had no effect on NADPH-CYP reductase activity (data not shown). Together with the results of kinetic analysis, this datum suggested that neo B interfered with the electron transfer from NADPH to CYP-substrate complex without directly affecting NADPH-CYP reductase reduction activity. In summary, our results indicated that lignans isolated from *J. p.* had the ability to affect CYP-catalyzed oxidations *in vitro*. To better understand the pharmacological implications and pharmaceutical applications, further studies should be done on the *in vivo* effects of *J. p.* extracts and lignans.

ACKNOWLEDGEMENTS

This work was supported by National Research Institute of Chinese Medicine and grant NSC89-2320-B077-001 from the National Science Council, R. O. C.

REFERENCES

- Guengerich, F. P. 1988. Roles of cytochrome P-450 enzymes in chemical carcinogenesis and cancer chemotherapy. *Cancer Res.* 48: 2946-2954.
- Nebert, D. W. 1989. The Ah locus; genetic differences in toxicity, cancer, mutation, and birth defects. *Crit. Rev. Toxicol.* 20: 153-174.
- Collins, J. F., Brown, J. P., Dawson, S. V. and Marty, M. A. 1991. Risk assessment for benzo(a)pyrene. *Regul. Toxicol. Pharmacol.* 13: 170-184.
- Levin, W., Wood, A., Chang, R., Ryan, D. and Thomas, P. 1982. Oxidative metabolism of polycyclic aromatic hydrocarbons to ultimate carcinogens. *Drug Metab. Rev.* 13: 555-580.
- Castonguay, A. 1992. Methods and strategies in lung cancer control. *Cancer Res.* 52 (9 Suppl): 2641s-2651s.
- Cassady, J. M., Baird, W. M. and Chang, C. J. 1990. Natural products as a source of potential cancer chemotherapeutic and chemopreventive agents. *J. Nat. Prod.* 53: 23-41.
- Fukamiya, N. and Lee, K. H. 1986. Antitumor agents, 81.¹ Justicidin-A and diphyllin, two cytotoxic principles from *Justicia procumbens*. *J. Nat. Prod.* 49: 348-350.
- Asano, J., Chiba, K., Tada, M. and Yoshli, T. 1996. Antiviral activity of lignans and their glycosides from *Justicia procumbens*. *Phytochemistry* 42: 713-717.
- Ward, R. S. 1993. Lignans, neolignans, and related compounds. *Nat. Prod. Reports* 10: 183-205.
- Chen, C. C., Hsin, W. C., Ko, F. N., Huang, Y. L., Ou, J. C. and Teng, C. M. 1996. Antiplatelet arylnaphthalide lignans from *Justicia procumbens*. *J. Nat. Prod.* 59: 1149-1150.
- Ueng, Y. F., Chang, Y. L., Oda, Y., Park, S. S. and Chen, C. F. 1999. *In vitro* and *in vivo* effects of naringin on cytochrome P450-dependent monooxygenase in mouse liver. *Life Sci.* 64: 2591-2602.
- Omura, T. and Sato, R. 1964. The carbon monoxide-binding pigment of liver microsomes. I. Evidence for its heme protein nature. *J. Biol. Chem.* 239: 2370-2379.
- Phillips, A. H. and Langdon, R. G. 1962. Hepatic triphosphopyridine nucleotide-cytochrome *c* reductase: Isolation, characterization, and kinetic studies. *J. Biol. Chem.* 237: 2652-2660.
- Nebert, D. W. and Gelboin, H. V. 1968. Substrate-inducible microsomal aryl hydroxylase in mammalian cell culture. I. Assembly and properties of induced enzyme. *J. Biol. Chem.* 243: 6242-6249.
- Pohl, R. J. and Fouts, J. R. 1980. A rapid method for assaying the metabolism of 7-ethoxyresorufin by microsomal subcellular fractions. *Anal. Biochem.* 107: 150-155.
- Brian, W. R., Sari, M. A., Iwasaki, M., Shimada, T., Kaminsky, L. S. and Guengerich, F. P. 1990. Catalytic activities of human liver cytochrome P-450III_{A4} expressed in *Saccharomyces cerevisiae*. *Biochemistry* 29: 11280-11292.
- Lowry, O. H., Roseborough, N. J., Farr, A. L. and Randall, R. L. 1951. Protein measurement with the Folin phenol reagent. *J. Biol. Chem.* 193: 265-275.
- Wood, A. W., Levin, W., Lu, A. Y. H., Yagi, H., Hernandez, O., Jerina, D. W. and Conney, A. H. 1976. Metabolism of benzo(a)pyrene and benzo(a)pyrene derivatives to mutagenic products by highly purified hepatic microsomal enzymes. *J. Biol. Chem.* 251: 4882-4890.
- Bauer, E., Guo, Z., Ueng, Y. F., Bell, L. C., Zeldin, D. and Guengerich, F. P. 1995. Oxidation of benzo(a)pyrene by recombinant human cytochrome P450 enzymes. *Chem. Res. Toxicol.* 8: 136-142.
- Guengerich, F. P. 1995. Human cytochrome P450 enzymes. In "Cytochrome P450: Structure, Mechanism, and Biochemistry". 2nd ed. pp. 473-535. Ortiz de Montellano, P. R. ed. Plenum press, New York, U. S. A.
- Nerurkar, P. V., Park, S. S., Thomas, P. E., Nims, R. W. and Lubert, R. A. 1993. Methoxyresorufin and benzyloxyresorufin: substrates preferentially metabolized by cytochromes P450 1A2 and 2B, respectively, in the rat and mouse. *Biochem. Pharmacol.* 46: 933-943.
- Burke, M. D., Thompson, S., Weaver, R. J., Wolf, C. R. and Mayer, R. T. 1994. Cytochrome P450 specificities of alkoxyresorufin O-dealkylation in human and rat liver. *Biochem. Pharmacol.* 48: 923-936.
- Ghosal, A., Sadrieh, N., Reik, L., Levin, W. and Thomas, P. E. 1996. Induction of the male-specific cytochrome P450 3A2 in female rats by phenytoin. *Arch. Biochem. Biophys.* 332: 153-162.

爵床木質素對 benzo(a)pyrene 氫氧化活性之抑制作用

翁芸芳* 陳建志 陳介甫

國立中國醫藥研究所
台北市立農街二段 155-1 號

(收稿：June 26, 2000；接受：September 6, 2000)

摘 要

爵床在傳統中藥中被用為治療發燒及喉嚨痛之處方用藥。由爵床乙醇萃取物中分離純化得 neojustin A (neo A), neojustin B (neo B), justicidin A (jus A), justicidin B (jus B) 及 chinensinaphthol methyl ether (CME) 五種木質素 (lignan)。分析木質對大白鼠肝微粒體單氧酵素催化 benzo(a)pyrene 羥化 (AHH) 活性之影響，結果顯示 10 μM 木質素對此活性具 21% 至 57% 之抑制作用，其中以 neo B 具最強之抑制作用，其 IC_{50} 為 $6 \pm 1 \mu\text{M}$ 。這些木質素 (10 μM) 對鼠肝微粒體 7-ethoxyresorufin O-脫乙基作用活性造成 20% 至 48% 之抑制作用。對 7-methoxyresorufin O-脫甲基作用活性，neo B 造成較其他木質素強之抑制作用，10 μM neo B 使 7-methoxyresorufin O-脫甲基作用活性下降 70%。對羧酮 6 β -羥化活性造成 19% 至 74% 抑制作用，而 neo B 之抑制作用最小。酵素動力學分析 neo B 抑制 AHH 活性之作用，結果顯示 neo B 為一競爭性 (competitive) 及非競爭性 (noncompetitive) 之混合型抑制物，其 K_i 及 K_i 分別為 2 μM 及 11 μM 。分析 neo B 對 NADPH 濃度與 AHH 活性相關性之結果顯示 neo B 非競爭性 (uncompetitive) 抑制 NADPH 支持之 AHH 活性。這些結果顯示木質素可對鼠肝單氧酵素催化活性造成不同程度之抑制影響，其中 neo B 對 AHH 活性具較強抑制作用，neo B 可能可降低 benzo(a)pyrene 之氧化活化作用。

關鍵詞：爵床，木質素，benzo(a)pyrene 羥化作用